


# Coronavirus (COVID-19)

---

# Weekly Epidemiological Bulletin



**Northern Ireland**

## Summary - Up to week 4 (31 January 2021)

To week 4, there have been a total of 104,075 laboratory confirmed cases<sup>1</sup> of COVID-19, including 2311 registered COVID-19 deaths<sup>2</sup> in Northern Ireland.

### COVID-19 case epidemiology



- 104,075 laboratory confirmed cases (17.5% from HSC laboratories)
- 53.8% of total cases are female
- In week 4, those aged 85+ had the highest case rate (307.2 per 100,000 population; 10.1% positivity)
- In week 4, Armagh, Banbridge and Craigavon had the highest case rate (340.4 per 100,000 population; 19.2% positivity)

### Care home outbreaks (suspected and confirmed)



- 567 confirmed COVID-19 outbreaks reported in total; includes 8 reported in week 4
- Involving 330 care homes (68.3% of all Northern Ireland care homes)
- The highest proportion of outbreaks (71.4%) were reported from the Belfast Trust area

### Sentinel testing



- Testing started 27 April 2020
- Number of individuals tested in total: 815 (4.4% positivity)

<sup>1</sup> Virological reports and the National Testing Programme.

<sup>2</sup> NISRA; 2020-21 - up to 22 January 2021

## COVID centres



- Testing started 15 June 2020 (one COVID centre: BHSCT)
- Virology data from the COVID centre became available from 1 July 2020
- Number of individuals tested 15 - 30 June 2020: 182 (all negative)
- Number of individuals tested 1 July – 22 November 2020: 864 (1.6% positivity)

## Critical care surveillance



- 522 confirmed COVID-19 individuals
- The majority of reported critical care cases were male (70.7%)
- Median age of cases was 62 years (range 21 – 90 years)

## Schools Surveillance



- 1641 COVID-19 incidents reported in 866 schools (58.4% of all Northern Ireland schools)
- 50.6% of incidents have been single cases, 36.7% were clusters of 2-5 cases; 12.7% were clusters of 6 or more cases
- Since schools opened, the highest proportion of incidents (70%) have been reported in Belfast

## Mortality surveillance



- In week ending 22 January 2021, the proportion of COVID-19 deaths registered was 38.4%. From the 19 March 2020 to week ending 22 January 2021 the proportion was 15.1%

- Excess deaths were reported in 2020 in weeks 13-20, 22, 44 and 45 and in 2021 in week 2; mainly in those over 65 years old.

### Testing surveillance virology



- Number of individuals tested in total: 993,098 (10.5% positivity)
- Number of individuals tested in;
  - HSC laboratories: 291,261 (29.3% of total tests)
  - National Testing Programme: 701,837 (70.7% of total tests)

## Introduction

COVID-19 is a new illness that can affect your lungs and airways. It's caused by a type of virus called SARS-CoV2 (coronavirus).

The Public Health Agency (PHA) Health Protection team has developed this report with the primary focus of looking at the demographic characteristics (age, sex and geographical location) of people affected by the virus. It also looks at some of the wider impact of the virus on the healthcare system, comparing recent trends in activity with historic norms.

There is a large amount of data being regularly published regarding COVID-19 (for example, the Department of Health Dashboard and *Deaths involving coronavirus in Northern Ireland* by the Northern Ireland Statistics and Research Agency). This report presents data from existing and newly developed PHA Health Protection surveillance systems that monitor COVID-19 activity in Northern Ireland and complements the range of existing data currently available.

As this is an emerging pandemic the systems used will constantly evolve and the complexity of the analysis will increase. All updates will be documented in “what’s new” section below.

Unless otherwise stated, data is presented using epidemiological weeks (a standardised method of counting weeks [Monday-Sunday] to allow for the comparison of data year after year). This is dependent on the data available and annual comparisons are not yet possible as the virus only emerged in 2020.

The data included in this report is the most up to date data available at the time of the report; however this is subject to change as the data is subject to ongoing quality assurance.

There is a large amount of data being regularly published regarding COVID-19 (for example, [Department of Health COVID-19 Daily Dashboard Updates](#) and [NISRA Deaths Registered Dashboard](#)). This bulletin complements the range of existing data currently available.

## Contact tracing

Contact tracing is the process of identifying, assessing, and managing people who have been exposed to a disease to prevent onward transmission ([WHO](#)). Contact tracing can help break the chains of transmission of COVID-19 and is an essential public health tool for controlling the virus.

Contact tracing seeks to limit and prevent the spread of infections such as COVID-19. It works by identifying a confirmed case and asking them who they have been in contact with. Individual contacts are considered high risk if they have spent more than 15 minutes in close contact with a confirmed case without personal protection. This means that those who have casually passed by someone on the street will not be considered high risk. The person with a confirmed infection and their close contacts will be given advice regarding symptom management and the need to self-isolate to prevent wider spread of the virus. This advice is based on information available on the PHA [website](#) and includes social distancing, handwashing and cleaning in the home to help protect people who are at risk. We can also advise people on how to best look after those in their care.

The most up-to-date contact tracing management service update (issued 28 January 2021) can be found [here\\*](#).

The StopCOVID NI contact tracing app is now [available](#) from the Google or Apple App store.

\*These are experimental performance and activity data and provide a snapshot of contact tracer activity. Data reported relates to a live operational system which includes case and contact activity in progress or in a queue. It is based on manually recorded information and data extracted from current contact tracing systems and reporting methods and parameters may change over time.

Automatic reporting in future may create a discontinuity in figures. New IT systems and data outputs often take some time to bed in. Data should therefore be treated with caution while the system and understanding of the data develops. At this stage, there is a risk of data entry errors or delay, which may require that data are revised and updated in future. The process of finding and removing duplicate records may also need refining, which could result in revisions to the data.

## Clusters

**Definition:** A cluster is currently defined as two or more laboratory confirmed cases of COVID-19 among individuals associated with a key setting, who have illness onset dates within a 14 day period. Key settings in which clusters have occurred include: workplaces, retail, hospitality and leisure premises as well as educational settings<sup>3</sup>.

### Comment:

Number of all clusters (open and closed) that have been recorded by the contact tracing service up to 5pm Monday 1 February 2021. Note: the reporting period for cluster data is slightly different to the remainder of the report in order to provide the most up to date cluster information at the time of the bulletin.

There have been 104 new clusters in the seven days up to 1 February 2021<sup>4,5</sup>. From week to week the number of clusters may change due to ongoing updates to the source information following detailed risk assessments. For this reason, we would discourage making direct comparisons between the cumulative number of clusters reported each week, with the number reported in the current week the most accurate at the time of the report.

In total, up to 1 February 2021, a total of 379 clusters with greater than five people have been identified in the following council areas; Antrim and Newtownabbey (n=36), Ards and North Down (n=11), Armagh City, Banbridge and Craigavon (n=44), Belfast City (n=82), Causeway Coast and Glens (n=14), Derry City and Strabane (n=37), Fermanagh and Omagh (n=23), Lisburn and Castlereagh City (n=19), Mid and East

---

<sup>3</sup> COVID-19 transmission is most common in household settings. The number of affected households is not reported.

<sup>4</sup> Note: the reporting period for cluster data is slightly different to the remainder of the report in order to provide the most up to date cluster information at the time of the bulletin. Some clusters may overlap (larger clusters may contain or overlap with several smaller clusters).


<sup>5</sup> From week to week the number of clusters may change due to ongoing updates to the source information following detailed risk assessments. For this reason, we would discourage making direct comparisons between the cumulative number of clusters reported each week, with the number reported in the current week the most accurate at the time of the report.

Antrim(n=34), Mid Ulster (n=46) and Newry, Mourne and Down (n=33). In addition, there have been 1365 clusters across Northern Ireland with fewer than five people.

*Source: Contact Tracing Service / PHA Health Protection Service*




## Case epidemiology




**Figure 1. Laboratory confirmed COVID-19 cases by epidemiological week and source (HSC Laboratory testing and the National Testing Programme), 2020-21**

Figure 1 represents the number of new weekly cases reported to the PHA (bars) and the cumulative number of cases (dashed line). Reporting is likely to be incomplete for the most recent week due to natural delays in samples reaching the labs, being tested and the information being reported.


From the end of September (week 40) we have seen a large increase in cases and increasing cumulative confirmed cases, peaking during week 42. This is mainly due to increasing clusters, increasing community transmission and contact tracing within a variety of settings. Although there were decreasing cases from the end of October (week 43) to week 48, from week 49 to week 53 there were increasing weekly cases. There has now been a decrease in weekly cases since week 53.




**Figure 2. Laboratory confirmed cases, by age, sex and source (HSC Laboratory testing and the National Testing Programme), 2020-21**



**Figure 3. Laboratory confirmed cases per 100,000 population, by age and sex, for all testing data combined, 2020-21**



**Figure 4. Laboratory confirmed cases, by age, sex and source (HSC Laboratory testing and the National Testing Programme), for weeks 3 and 4**



**Figure 5. Laboratory confirmed cases per 100,000 population, by age and sex, for all testing data combined, for weeks 3 and 4**

Figures 2 and 3 represents the cumulative number of cases reported by HSC laboratories and the National Testing Programme, and overall case rates per 100,000 population, respectively. HSC laboratory cases were mainly detected at the beginning of the pandemic in hospital settings, resulting in higher cases and rates among the older age groups. With the introduction of the National Testing Programme cases it has become the main source of case data as a result of enhanced community testing enabling us to detect a greater spectrum of disease. From this data we have seen a higher number of cases among the 20-29 age group.

Figures 4 and 5 show similar to the cumulative numbers but restricted to the previous two epidemiological weeks. These show how the age groups of cases in the most recent weeks differ from the overall cumulative cases presented in figures 2 and 3. Also, more cases are being detected outside of hospital settings as part of the National Testing Programme.

Table 1. Total laboratory confirmed COVID-19 cases, by sex, for all testing data combined			
Age group	Sex		Total*
	Male	Female	
0 - 9	2015	1927	3942
10 - 19	4650	5338	9988
20 - 29	9106	10770	19876
30 - 39	8181	9692	17873
40 - 49	7193	8472	15665
50 - 59	7462	8605	16067
60 - 69	4511	4413	8924
70 - 79	2675	2734	5409
80+	2322	3984	6306
Unknown		2	2
<b>Total</b>	<b>48115</b>	<b>55937</b>	<b>104052</b>

\*Unknown sex for twenty three cases, these are not included in the total figures

Table 2. Laboratory confirmed COVID-19 cases, by Trust				
Trust Area	Epidemiological Week			
	2	3	4	Total
Belfast	241	194	117	5298
Northern	197	136	89	3759
South Eastern	77	85	86	2316
Southern	178	121	164	2825
Western	71	39	38	1431
Other*	5628	4137	3036	88363
Unknown	1	1	0	83
<b>Northern Ireland</b>	<b>6393</b>	<b>4713</b>	<b>3530</b>	<b>104075</b>

\*Other cases includes those from the National Testing Programme, NIAS, private nursing home residents, pathology services, GPs and hospices






Figure 6. Weekly laboratory confirmed case rates per 100,000 population, by age group, for all testing data combined, 2020-21

The case rates decreased in week 4 compared to the previous week in all age groups. The highest case rates were seen in the 85+ age group (307.2 per 100,000), which is lower than the peak of 970.1 per 100,000 in the 15-44 age group in week 53 (27 December 2020-03 January 2021). In week 4, positivity was highest in the 5-14 age group (18.0%). The lowest positivity was observed in the 0-4 age group (9.3%).




**Figure 7. Positivity (%) of laboratory confirmed COVID-19 cases by epidemiological week, overall and by sex (HSC Laboratory testing), 2020-21**



**Figure 8. Positivity (%) of laboratory confirmed COVID-19 cases by epidemiological week, overall and by sex (National Testing programme), 2020-21**






**Figure 9. Total laboratory confirmed cases, by Local Government District (LGD) and source (HSC Laboratory testing and the National Testing Programme), 2020-21**

Public Health Agency

Density map of confirmed COVID-19 cases during the week ending 31 January 2021





Source: Health Protection Department PHA  
 Produced on behalf of Health Intelligence PHA (02/02/2021)  
 Reproduced with the permission of Land and Property Services under delegated authority from  
 the Controller of Her Majesty's Stationary Office. © Crown copyright and database rights NIMA ES&LA210.4

From data extracted on 2nd February 2021


**Figure 10. Density map of confirmed COVID-19 cases, for all testing combined, 2020-21**

Figure 10 shows a contour density map based on the number of confirmed COVID-19 cases in week 4. The contour lines on the map indicate increasing density of cases, with the darkest shade of blue indicating where there is the greatest density of cases. The map highlights an increased number of areas with the highest density of COVID19 cases in week 4 and spread of the virus to other areas. However, information should be interpreted with caution as identified rates are based on testing which is not evenly spread across the region.




**Figure 11. Total laboratory confirmed cases per 100,000 population, by Local Government District (LGD), for all testing data combined, week 4 (25 - 31 January 2021)**


### Antrim and Newtownabbey




### Ards and North Down




### Armagh, Banbridge and Craigavon




### Belfast




### Causeway Coast and Glens




### Derry and Strabane



### Fermanagh and Omagh



### Lisburn and Castlereagh









Figure 12. Weekly laboratory confirmed cases per 100,000 population and proportion positive, by Local Government District (LGD) and Northern Ireland, for all testing data combined, 2020-2021

The case rates decreased in all areas in week 4 compared to week 3. Armagh, Banbridge and Craigavon had the highest rate in week 4 compared to other Local Government Districts (340.4 per 100,000 population). The overall Northern Ireland rate decreased from 248.9 to 186.4 per 100,000 population between weeks 3 and 4.

The highest positivity occurred in Armagh, Banbridge and Craigavon (19.2%). Northern Ireland's proportion positive in week 4 was 12.1%, a decrease from 15.8% in week 3. This is lower than the peak positivity of 26.4% reported across Northern Ireland in week 53 (27 December 2020 – 03 January 2021).

*Source: HSC Trust laboratory reports and the National Testing Programme*

## Deprivation

An analysis of COVID-19 related health inequalities relating positive test cases and COVID-19 related admissions between the most and least deprived areas of Northern Ireland, including variations across age, sex and urban and rural areas was [published](#) by Department of Health on 16 December 2020.

## Care home outbreaks





Figure 13. Confirmed COVID-19 care home outbreaks in Northern Ireland, 2020-21




**Figure 14. Proportion of care homes with confirmed or suspected COVID-19 in Northern Ireland by Trust, 2020-21**

Table 3. Proportion of care homes with confirmed COVID-19 outbreaks in Northern Ireland, by Trust			
Trust Area	Cumulative total of care homes with outbreaks in 2020	% of care homes with outbreaks	Total number of care homes
<b>Belfast</b>	65	71.4%	91
<b>Northern</b>	93	70.5%	132
<b>South Eastern</b>	68	61.3%	111
<b>Southern</b>	57	73.1%	78
<b>Western</b>	47	66.2%	71
<b>Northern Ireland</b>	330	68.3%	483

To week 4, a total of 567 confirmed COVID-19 care home outbreaks were reported, involving 330 care homes (68.3% of all Northern Ireland care homes). The highest proportion of care homes with confirmed COVID-19 outbreaks (71.4%) were reported from the Belfast Trust area.





**Figure 15. Number of care homes with a confirmed active symptomatic or asymptomatic COVID-19 outbreak<sup>6</sup> in Northern Ireland, 2020-21**

*Source: PHA Health Protection duty room reports from care homes*

<sup>6</sup> PHA began recording confirmed Covid-19 outbreaks as either symptomatic or asymptomatic on 1 August 2020. This means the numbers represented on the graph may not equal the total active confirmed COVID-19 outbreaks. Confirmed COVID-19 outbreaks reported prior to 1 August 2020 and are still ongoing are not included in this graph. Additionally, other respiratory outbreaks are not included.

## Sentinel testing

Table 4. COVID-19 activity in Northern Ireland Sentinel GP Practices*, week 4, 2021			
Period	Individuals tested	Number positive	Proportion positive
Current week	20	1	5.0%
<b>Total</b>	<b>815</b>	<b>36</b>	<b>4.4%</b>

\*Sentinel testing started 27 April 2020; excludes care home residents and healthcare workers


## COVID centre testing

Table 5. COVID-19 activity in Northern Ireland COVID Centres*, week 4, 2021			
Period	Individuals tested	Number positive	Proportion positive
Current week	-	-	N/A
<b>Total</b>	<b>864</b>	<b>14</b>	<b>1.6%</b>

\* One COVID centre operational from 15 June 2020 (BHSC); virology data in table above from 01 July 2020. Data provided from the COVID centre directly reported 182 individuals tested between 15 June and 30 June 2020 inclusive. All results were negative. This data is subject to change as we continue to quality assure the COVID centre information against virology.

Source: HSC Trust laboratory reports and the National Testing Programme

## Critical care surveillance



Change in surveillance procedure\*

\*Since start of week 40 (28 September 2020), data collection for critical care surveillance has been streamlined to coincide with the well-established surveillance of influenza patients in critical care in conjunction with the Critical Care

Figure 16. ICU/HDU COVID-19 cases by sample result week, 2020-21




Figure 17. ICU/HDU COVID-19 cases, by age and sex, 2020-21

To week 4, there have been 522 individuals admitted to critical care with confirmed SARS-CoV2 reported to the PHA. Week 2 had the highest number of ICU reports with a positive result (n=49).

Of the 522 individuals, 70.7% (n=369) were male. The ages ranged from 21 years to 90 years, with a median age of 62 years.

*Source: PHA COVID-19 critical care surveillance online reporting system and the Critical Care Network Northern Ireland (CaNNI)*

The Intensive Care National Audit and Research Centre (ICNARC) publish a report on patients critically ill with COVID-19 (<https://www.icnarc.org/Our-Audit/Audits/Cmp/Reports>). There is also a specific report which can be downloaded presenting analysis of data on patients critically ill with confirmed COVID-19 reported to ICNARC up to 23:59 on 28 January 2021 from critical care units in Northern Ireland participating in the Case Mix Programme (the national clinical audit for adult critical care).

## Schools Surveillance

Information on school COVID incidents is based on situations reported to PHA COVID School Team.

These include:

- **Single confirmed case** of COVID-19 (SARS-CoV-2) in a student or member of staff in the school setting.

The incident is closed after 14 days if there have been no further cases.

- **Cluster of two or more confirmed cases of COVID** (SARS-CoV-2) in a student or member of staff in the school setting within a 14 day period.

The incident is closed after 14 days if there have been no further cases from the last case

The PHA COVID School Team carries out contact tracing of cases that attend a school in collaboration with PHA Test and Trace Programme. Clusters are also further investigated by the School Team in liaison with local partners.

Data is collected on the number of COVID school incidents reported to the PHA COVID School Team since schools reopened.

The number of cases that have been reported by schools to the PHA school team is also included in this section to provide high level information on cases broken down by pupil and staff status. It is important to note that the definitive source for the number of COVID-19 confirmed cases in school aged children is from the HSC Laboratory testing and the National Testing Programme. Direct comparisons should not be made with laboratory data as the school teams figures may underestimate of laboratory data.

Table 6 shows the number of school incidents by type of school that have been reported to the PHA School team up to the end of week 4.

The figures are a snapshot of incidents recorded at the time of data extraction. A school may have had more than one incident since opening. Figures should not be compared from week to week as the number will include new reports and further cases of existing incidents.

<b>Table 6. Number of COVID-19 Incidents in Schools</b>	
<b>School Type</b>	<b>Total to date</b>
<b>Preschool</b>	106 (6.5%)
<b>Primary</b>	938 (57.2%)
<b>Post Primary</b>	495 (30.2%)
<b>Special</b>	102 (6.2%)
<b>Total</b>	1641

Table 7 shows the number of school incidents by type of school and also type of incident i.e. single case in a 14 day period or cluster of cases within a 14 day period.

Clusters have been further broken down into those with 2 to 5 cases and 6 or more cases.

Table 7. Number of Incidents by School and Incident Type			
Incident Type	School Type	Total to date	Proportion
Single Case	Preschool	77	9.3%
	Primary	541	65.1%
	Post Primary	162	19.5%
	Special	51	6.1%
	All	831	
Cluster (2-5 cases)	Preschool	26	4.3%
	Primary	342	56.8%
	Post Primary	192	31.9%
	Special	42	7.0%
	All	602	
Cluster (>5 cases)	Preschool	3	1.4%
	Primary	55	26.4%
	Post Primary	141	67.8%
	Special	9	4.3%
	All	208	

## Cumulative number of schools affected by at least one case of COVID-19

A school may have had more than one incident since opening on 24th August. Table 8 shows the cumulative number of schools that have had at least one school incident up to the end of week 4.

The 1641 school incidents have occurred in 866 schools in Northern Ireland. Overall 58.4% of schools have had at least one COVID-19 case in a pupil or member of staff.

Table 8. Number of Schools with a COVID-19 Incident			
School Type	No. Schools that have had at least one case	Total number of schools in Northern Ireland	Proportion of school in Northern Ireland that have had at least one case
Preschool	88	458	19.2%
Primary	554	792	69.9%
Post Primary	190	194	97.9%
Special	34	39	87.2%
<b>Total</b>	<b>866</b>	<b>1483</b>	<b>58.4%</b>



### Trend of school incidents

The following information includes the number of incidents in schools since they first reopened until the end of week 4 (31 January 2021).





Figure 18. Number of COVID-19 incidents in schools, by school type, week 33 - 4

### Cumulative School Incidents by Local Government District

The following information includes the cumulative number of incidents in schools by LGD since they first reopened until the end of week 4 (31 January 2021).

The cumulative community rate per 100,000 population is also shown in the figure.



**Figure 19. Proportion of schools with a COVID-19 incident and overall background community rate by 100,000 (2020) by Local Government District (LGD)**

## Cumulative number of COVID-19 cases reported by schools to PHA School Team

Since schools opening on 24th August until end of week 4, there have been 5,033 confirmed COVID-19 cases that occurred at any point during this time reported by schools to the PHA School Team.


The definitive source for the number of COVID-19 confirmed cases in school aged children is from the HSC Laboratory testing and the National Testing Programme. Direct comparisons should not be made with laboratory data.

Staff member includes teaching and non-teaching staff.

Table 9. Number of COVID-19 cases reported by schools where information is available on pupil / status, up to week 4				
School	Pupil Case	Staff Case	Total	Proportion of all cases that are pupils
Preschool	67	104	171	39.2%
Primary	1051	886	1937	54.3%
Post Primary	2075	605	2680	77.4%
Special	67	178	245	27.3%
All	3260	1773	5033	64.8%

Table 10. Number of COVID-19 cases in school aged children reported by schools where information is available as a proportion of all school age children, up to week 4		
School Type	Pupil cases	Proportion of all school aged pupils in Northern Ireland
Preschool	67	0.02%
Primary	1051	0.31%
Post Primary	2075	0.61%
Special	67	0.02%
All	3260	0.95%

Source: PHA COVID-19 Schools Team, Department of Education school statistics



**Figure 20. Weekly COVID-19 laboratory confirmed case rates per 100,000 population, by age group, for all testing data combined, in those aged 21 and under, 2020-21**

All of the age groups decreased in week 4 compared to week 3. The 19-21 age group had the highest case rates (225.8 per 100,000), followed by the 17-18 age group (184.9 per 100,000).

*Source: HSC Trust laboratory reports and the National Testing Programme*

## Mortality surveillance

### Medical Certificate of Cause of Death for confirmed / suspected COVID-19

The Northern Ireland Statistics and Research Agency (NISRA) provide the weekly number of **registered respiratory and COVID-19 deaths each Friday ([here](#))**. In week ending 22 January 2021, the proportion of COVID-19 deaths registered was 38.4%, and from 19 March 2020 to week ending 22 January 2021 the proportion of COVID-19 deaths registered was 15.1%.

Weekly published data are provisional and is based on registrations of deaths, not occurrences. The majority of deaths are registered within five days in Northern Ireland. Respiratory deaths include any death where terms directly relating to respiratory causes were mentioned anywhere on the death certificate (this includes Covid-19 deaths). This is not directly comparable to the ONS figures relating to 'deaths where the underlying cause was respiratory disease'. Figures relate to all deaths registered up to 22 January 2021 with a mention of COVID on the death certificate. Please note: Where COVID is mentioned in part 1 it may not be the underlying cause of death. Covid-19 deaths include any death where Coronavirus or Covid-19 (suspected or confirmed) was mentioned anywhere on the death certificate. NISRA quarterly statistics provide detail of underlying cause following coding to ICD-10 rules; figures are available [here](#).





Figure 21. NISRA registered COVID-19 deaths by sex and age group, up to week ending 22nd January 2021



Up to week 52, the Average deaths (previous 5 years) period ranged from 2015-2019. From 2021 week 1 onwards, this period ranges from 2016-2020

Figure 22. Northern Ireland registered deaths, including COVID-19 associated deaths, Week 11 (ending 20 March 2020) to Week 3 (ending 22 January 2021)

**Table 11. Northern Ireland registered deaths, including COVID-19 associated deaths, Week 11 (ending 20 March 2020) to Week 3 (ending 22 January 2021)**


Registration week (ending Friday)	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	1	2	3
COVID-19 death certificate, Part 1	0	8	50	75	100	123	120	75	65	45	35	18	15	14	11	10	7	0	6	1	4	3	4	3	2	5	4	7	1	6	15	36	50	77	82	89	74	86	73	76	72	82	122	136	152
COVID-19 death certificate, Part 2	1	1	5	1	1	5	4	9	9	8	14	2	6	3	1	1	2	2	1	0	1	1	2	1	1	2	4	2	1	5	2	6	1	1	14	11	7	12	14	6	16	11	23	17	30
Non-COVID-19 deaths	270	278	379	359	323	342	303	252	322	272	267	284	271	273	283	278	266	238	300	272	275	274	307	299	231	289	314	314	326	337	261	349	317	304	310	296	267	289	279	268	222	240	423	290	292
Average deaths (previous 5 years)	297	320	298	295	290	284	320	274	295	290	279	271	293	286	270	288	255	264	265	282	276	265	263	259	255	276	282	288	286	300	295	286	284	296	297	319	311	322	322	344	281	280	417	399	375
Total deaths (2020P)	271	287	434	435	424	470	427	336	396	325	316	304	292	290	295	289	275	240	307	273	280	278	313	303	234	296	322	323	328	348	278	391	368	386	406	396	348	387	366	350	310	333	568	443	474

Up to week 52, the Average deaths (previous 5 years) period ranged from 2015-2019. From 2021 week 1 onwards, this period ranges from 2016-2020

Source: Northern Ireland Statistical Research Agency (NISRA)



## All-cause excess deaths




**Figure 23. Weekly observed and expected number of all-cause deaths in all ages, week 40 2017 - week 4 2021**

In 2020, excess all-cause deaths were reported in epidemiological weeks 13 to 20, week 22 and weeks 44 and 45. During 2021, excess deaths were reported in week 2 (11th – 17th January). This increase in deaths happened outside the influenza season and at a time when we know flu was not circulating ([here](#)).

While these more recent excesses have occurred within the flu season, reports show flu was not widely circulating. This suggests the excess mortality may in part be related to COVID-19 deaths and to a fall in presentation to hospital with other conditions (data not shown). Excess deaths were mainly in those over 65 years, which is in line with the age profile of COVID-19 deaths. Despite delay correction, reported mortality data are still provisional due to the time delay in registration and observations which can vary from week to week; not all registrations for the current week will have been included this bulletin.

## Virology testing surveillance



Total individuals tested include those that were reported as indeterminate

**Figure 24. Weekly number of individuals tested for SARS-CoV2 and proportion positive, by source (HSC Laboratory testing and the National Testing Programme), 2020-21**

**Table 12. COVID-19 activity in Northern Ireland, for all testing data combined, week 4, 2021**

Period	Individuals tested	Number positive	Proportion positive
Current week	29087	3530	12.1%
<b>Total</b>	<b>993,098</b>	<b>104,075</b>	<b>10.5%</b>

**Table 13. COVID-19 activity in Northern Ireland (HSC laboratory), week 4, 2021**

Period	Individuals tested	Number positive	Proportion positive
Current week	6853	532	7.8%
<b>Total</b>	<b>291,261</b>	<b>18,169</b>	<b>6.2%</b>

**Table 14. COVID-19 activity in Northern Ireland (National Testing Programme), week 4, 2021**

Period	Individuals tested	Number positive	Proportion positive
Current week	22234	2998	13.5%
<b>Total</b>	<b>701,837</b>	<b>85,906</b>	<b>12.2%</b>

*Source: HSC Trust laboratory reports and the National Testing Programme*

To week 4, the total number of individuals tested was 993,098, positivity 10.5%. Overall, more individuals have now been tested as part of the National Testing Programme, and positivity is now higher (12.2%) compared to HSC laboratories (6.2%).

## Global situation

As of 3 February, [WHO](#) has been notified of 103,201,340 confirmed cases of COVID-19, including 2,237,636 related deaths.

## Appendix

### PHA Health Protection COVID-19 surveillance systems

The PHA Health Protection Directorate has established the following surveillance systems to monitor COVID-19 activity across the spectrum of community and health care settings. As new systems are developed they will be added to this report.

### Case epidemiology

SARS-CoV2 testing was first developed by the National Reference Laboratory (Public Health England) for all of the United Kingdom on 24 January 2020. On 7 February 2020, SARS-CoV2 testing was developed locally by the Regional Virus Laboratory, Belfast Health and Social Care (HSC) Trust and performed testing across Northern Ireland. Since 23 March, 28 March, 3 April and 13 May respectively, Northern HSC Trust, Southern HSC Trust, Western HSC and South Eastern HSC Trust laboratories, have been performing SARS-CoV2 testing.

The PHA Health Protection Directorate laboratory surveillance system collates SARS-CoV2 laboratory data on all tests from HSC Trust laboratories.

As an individual may have more than one test for clinical purposes, the laboratory data is then collated to enable monitoring of individuals rather than tests performed by laboratories. This is done using the Organism-Patient-Illness-Episode (OPIE) principle, a standard approach used across the UK.<sup>7</sup> The episode length used nationally is 6 weeks (42 days), and is being reviewed as more data becomes available.

---

<sup>7</sup> Public Health England. 2016. Laboratory reporting to Public Health England: A guide for diagnostic laboratories. [ONLINE] Available at:

If an individual is infected on two separate occasions by the same organism (within the episode of infection) they will be represented by one distinct record. It is still unclear to what extent second infections occur in COVID-19. The exception to this is where the first result is negative and is then followed by a positive result on a second occasion. In such circumstances, the later positive result will be recorded rather than the earlier negative one. If an individual is infected on two separate occasions by the same organism (outside the episode of infection with recovery implied) they will be represented by two distinct records, regardless of the test result. This is a standard approach which is taken across a range of infectious diseases.

All laboratories report a standardised data set which includes individual demographics, test result and source (location) at the time the specimen was taken. Data is collated to produce information on the number and trend of individuals tested at HSC Trust laboratories and the number and trend of confirmed cases in Northern Ireland.

### National Testing Programme

The National Testing Programme in Northern Ireland consists of drive through (regional test sites), mobile test unit sites, home testing and satellite testing of nursing homes.

Everyone in Northern Ireland with symptoms of coronavirus is now eligible for testing.

Testing is prioritised through the website gov.uk for essential workers who are self-isolating because they are symptomatic, or have household members who are symptomatic, to help enable essential workers to return to work as soon as safe.

Testing is available for the general public through the website nhs.uk.

The StopCOVID NI contact tracing app is now [available](#) from the Google or Apple App store.

Testing for non-HSC essential workers and the general public is currently conducted in drive-through sites operating in Belfast, Enniskillen, Derry/Londonderry and Portadown. In addition there is a mobile testing unit currently operating within Northern Ireland.

Home testing can be requested by any individual meeting the criteria with a test kit(s) being mailed to the individual and household contacts.

Tests are processed in laboratories outside the normal health and social care network and data fed back to the Public Health Agency via the Business Services Organisation.

The data has been included in the case epidemiology and virology testing surveillance sections. This data should be interpreted with caution, when interpreted alongside the HSC laboratory data, because it includes testing undertaken as part of the outbreak response i.e. possibly asymptomatic people with a certain age, gender or area profile. Testing numbers may be skewed to different local government districts depending on whether an outbreak was detected and managed.

For more information see [here](#).

### Care home outbreak surveillance

A care home is a term that includes all nursing homes and residential homes in Northern Ireland that are registered with the Regulation and Quality Improvement Agency (RQIA) and can either be HSC Trust or independent sector owned. There are 472 active care homes in Northern Ireland.

All care homes have a requirement to notify the PHA Health Protection duty room of suspected outbreaks of any infectious disease. A suspected outbreak of COVID-19 occurs when two or more residents and/or staff meet the case definitions for suspected COVID-19, confirmed COVID-19, influenza-like illness or worsening shortness of breath.

The PHA Health Protection Directorate care home outbreak surveillance system collects and collates data on all initial notifications of suspected COVID-19 outbreaks from the duty room clinical records.

The care home COVID-19 outbreak surveillance system is updated every day to reflect public health management. If the risk assessment subsequently excludes an outbreak of the initial notification then the surveillance data will be updated.

Currently, care homes with multiple facilities, i.e. nursing and residential, but the same name may be reported as one outbreak, rather than two (if both units are affected) which may underestimate the number of care homes affected.

## Primary care surveillance

### a. Sentinel testing

The GP sentinel testing surveillance system builds on the existing flu sentinel testing system where 36 general practices ('spotter' practices), representing approximately 11% of practices across Northern Ireland, are commissioned to carry out flu testing in suspected influenza-like illness.

Individuals registered at a spotter practice with symptoms of suspected COVID-19 and who are well enough to self-care in their own home are referred to a Trust testing facility for testing. The service commenced in 13 spotter practices in Belfast and South Eastern HSC Trust locality at the end of April and is currently being rolled out to the other 23 practices in Northern, Southern and Western HSC Trust localities.

Laboratories reports from spotter practices are identified from the laboratory (virology) surveillance and are collated to produce information on the number of individuals tested and the number of confirmed cases.

### b. COVID centre testing

A COVID centre is a separate facility created as an extension of primary care to help direct suspected COVID positive patients for assessment.

This keeps practices free to deal with any other medical problems. Triage will still occur at the practice, most likely via phone followed by referral to the centre if required.

There are three categories of patient that might be assessed at a COVID centre:

1. patients symptomatic for COVID, or already test positive who are clinically worsening: there will also be direct pathways for investigation and/or admission from the centre
2. patients where there is diagnostic uncertainty: symptoms similar to COVID but could be another clinical problem ranging from tonsillitis to meningitis requiring an assessment to exclude or confirm
3. patients being discharged from hospital: this group will grow with time but on many occasions will still have a need for clinical assessment and follow up.

Centres are staffed by GPs, helped by other members of staff, including nurses, health care workers etc.

Centres run from 8am to 10pm and see patients after triage and referral (by CCG) from the practice.

Patients can either be seen in their car outside the centre if a straightforward examination is needed, or brought into the centre for assessment. Patients are told to wait in their car until phoned to come in to prevent any crowding or grouping of patients.

Centres are hosted by the trusts and operate in each trust area.

### **Critical care surveillance**

Until 28<sup>th</sup> September 2020, the PHA Health Protection COVID-19 critical care online reporting system captured the incidence of COVID-19 infections in critical care and aims to improve the understanding of severe disease.

This system should complement critical care data collected by the Health and Social Care Board for service planning purposes and the publicly available reports on COVID-19 in critical care Northern Ireland by the Intensive Care National Audit and Research Centre (iCNARC) ([here](#)).



Since 28<sup>th</sup> September 2020, data collection for critical care surveillance has been streamlined to coincide with the well-established surveillance of influenza patients in critical care in conjunction with the Critical Care Network Northern Ireland (CaNNI).

Data is collected on all individuals admitted to an Intensive Care Unit (ICU) or High Dependency Unit (HDU) with a positive SARS-CoV2 result, from either before or during the ICU/HDU admission.

## Schools Surveillance

Information on school COVID incidents is based on situations reported to PHA COVID School Team. These include:

- **Single confirmed case** of COVID (SARS-CoV-2) in a student or member of staff in the school setting. The incident is closed after 14 days if there have been no further cases.
- **Cluster of two or more confirmed cases of COVID** in a student or member of staff in the school setting within a 14 day period. The incident is closed after 14 days if there have been no further cases from the last case

PHA COVID School Team carries out contact tracing of cases that attend a school in collaboration with PHA Test and Trace Programme. All clusters are also investigated by the School Team in liaison with local partners.

Data is collected on the number of COVID school incidents reported to the PHA COVID School Team since schools reopened. This is the first week of reporting school surveillance.

## Mortality surveillance

### Medical Certificate of Cause of Death for confirmed/suspected COVID-19

The traditional method for examining the number of deaths, and the range of causes of death, takes information from death certificates that are reported to the General Registrar's Office (GRO). The death certificate contains two parts. Part 1 describes the immediate causes of death and Part 2 provides information on related conditions that may

also have contributed to death. The numbers of deaths from COVID-19 are based on COVID-19 being recorded on any part of the death certificate (i.e. Part 1 or Part 2).

These include all deaths in which a doctor feels that COVID was either a direct or indirect cause of death. It includes confirmed cases (deaths with a positive laboratory result) and probable or suspected cases, where a doctor assesses that COVID was a cause of death but there is either no lab test or the test was negative. It captures deaths in all settings, such as hospitals, care homes, hospices and the community. It takes up to five days for most deaths to be certified by a doctor, registered and the data processed, meaning these deaths will be reported on about a week after they occurred.

Inclusion of references to COVID-19 in Part 2 of the death certificate may slightly over estimate the number of individuals where COVID-19 is a significant contributor to death.

### All-cause excess deaths

The PHA Health Protection Directorate reports the weekly number of excess deaths from any cause for Northern Ireland using the Mortality Monitoring in Europe (EuroMOMO) model. EuroMOMO provides a coordinated, timely and standardised approach to monitoring and analysing mortality data across the UK and Europe, to ensure that signals are comparable between countries. Further information is available [here](#).

Based on mortality data supplied by NISRA, EuroMOMO produces the number of expected and observed deaths every week, corrected for reporting delay and standardised for the population by age group and region. Excess mortality is reported if the number of observed deaths exceeds the number of expected deaths, and is defined as a statistically significant increase in the number of deaths reported over the expected number for a given point in time.

## Case definitions

Case definitions are determined by Public Health England, on the advice of the New and Emerging Respiratory Virus Threats Advisory Group (NERVTAG). As the pandemic evolves and more evidence emerges the definitions will change to ensure individuals are appropriately identified.

### Possible case of COVID-19 (as of 28 September 2020)

As of 2<sup>nd</sup> October, case definitions for inpatient and community settings were consolidated into one list. Unusual presentations are also highlighted.

Individuals with

- new continuous cough **OR**
- high temperature **OR**
- a loss of, or change in, normal sense of smell (anosmia) or taste (ageusia)

Individuals with any of the above symptoms but who are well enough to remain in the community should follow the [stay at home guidance](#) and [get tested](#).

Clinicians should be alert to the possibility of atypical presentations in patients who are immunocompromised.

Alternative clinical diagnoses and epidemiological risk factors should be considered.

A wide variety of clinical symptoms have been associated with COVID-19.

Patients with acute respiratory infection, influenza-like illness, clinical or radiological evidence of pneumonia, or acute worsening of underlying respiratory illness, or fever without another cause should have a SARS-CoV-2 test, whether presenting in primary or secondary care.

In addition, the following situation should prompt clinicians to consider SARS-CoV2 testing:

- Onset of delirium (acute confusion) in older people, or in those with dementia or cognitive impairment. New infections in people with dementia may be manifest as delirium.

## Confirmed case of COVID-19

An individual with clinical symptoms and a positive SARS-CoV2 specimen result.

## Critical care COVID-19 case

A case that has either been admitted to an ICU/HDU in Northern Ireland with a pre-existing positive result for SARS-CoV2, or received a positive result for SARS-CoV2 post-admission to ICU/HDU.

## Influenza-like Illness (ILI)

Acute respiratory disease with sudden onset of symptoms and:

- at least one systemic symptom (fever  $\geq 37.8^{\circ}\text{C}$ , myalgia, malaise, headache) AND
- at least one respiratory symptom: cough (with or without sputum), shortness of breath (and/or wheezing), sore throat, nasal discharge, sneezing or congestion

## Further Information

This bulletin is produced by the Health Protection Surveillance Team on behalf of the Director of Public Health. Correspondence should be directed to: Dr Stephen Bergin, Acting Director of Public Health, Public Health Agency, 12 – 22 Linenhall Street, Belfast, BT2 8BS. Email: [Stephen.Bergin@hscni.net](mailto:Stephen.Bergin@hscni.net) .

© Public Health Agency: 3 February 2021.