



Crossnacreevy Suction Trap 2020

W/E 14/08/20		Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Total
		08-Aug	09-Aug	10-Aug	11-Aug	12-Aug	13-Aug	14-Aug	
<i>Brachycaudus helichrysi</i>	(Leaf-curling plum aphid)								
<i>Macrosiphum euphorbiae</i>	(Potato aphid)								
<i>Metopolohium dirhodum</i>	(Rose-grain aphid)								
<i>Rhopalosiphum spp.</i>		10	21	19	37	11	10	21	129
<i>Sitobion avenae</i>	(Grain aphid)								
Others		1	1	9	40	20	1	1	73
Total		11	22	28	77	31	11	22	202
Notes		Humid and increasingly unsettled weather during this period, <i>Rhopalosiphum spp.</i> numbers continuing to increase. However, still below seasonal average.							



Weekly Cereal Aphid Monitor 2020

Cereal aphid counts week ending 14th August 2020



Headlines:

We are pleased to initiate suction trapping at Crossnacreevy 2020

- Complete weekly suction trapping week ending 14th August 2020
- 129 *Rhopalosiphum* spp. recorded in trap
- *Rhopalosiphum* spp. numbers are increasing, but remain significantly below the 13-year mean
- No *Sitobion avenae* adults recorded this week
- No *Metopolophium dirhodum*

How is BYDV transmitted?

The aphids transmit BYDV into winter cereals by two ways:

1. Cereal volunteers, or grass weeds within a field can act as hosts for aphids after stubble destruction. This type of infection typically causes large discrete patches of severely infected plants.
2. Winged aphids flying from grass or cereal volunteers elsewhere can also introduce the virus into the newly emerged crops during the autumn. In some regions, this is the most common route for BYDV infection.

Risk forecasting

The Aphid Migration graph provides an indication of aphid numbers in Northern Ireland and consequently the relative risk of BYDV. This data is collected and updated weekly by AFBI, Newforge Lane during the high-risk period. The current situation is displayed against a reference “12- year average” to indicate whether the aphid numbers are significantly higher than usual. This is used to assess the degree of risk of BYDV infection in conjunction with the emergence dates of winter cereal crops. The advice based on this assessment could be:

High risk

A large early migration of aphid vectors in conjunction with the coincidence of early emergence of crops.

Action: Application of insecticidal spray to crops which have reached the 2-leaf stage of growth. This should be followed up after 2 weeks with a crop inspection for further aphid arrivals.

Moderate risk

Migration at or slightly higher than the 12- year average and crops emerging later.

Action: Application of insecticidal spray to crops (2-leaf stage) emerged before the end of the aphid migration (usually early November).

Low risk

Migration below the 12-year average.

Action: Monitor crops regularly for presence of aphid vectors. Application of insecticidal spray only to those crops where aphids are found.

Aerial introduction of BYDV by migrating aphids results in point infections scattered throughout the crop. The real damage is caused when these initial aphids give rise to colonies which disperse and carry the infection to surrounding plants. Insecticidal sprays are used to limit this spread of BYDV so that the overall level of infection is insignificant when the crop matures.

BYDV control

The Aphid Migration graph provides an indication of aphid numbers in Northern Ireland and consequently the risk of BYDV, although this may vary regionally. During the high-risk autumn period this data is collected and updated weekly by AFBI, Newforge Lane. To view the updated reports and for more information click the link to weekly reports. (<https://www.afbini.gov.uk/publications/cereal-aphids-weekly-results-2019>)

In situations where there is a weedy stubble or a large number of volunteers cereal plants, this "green bridge" should be destroyed by desiccation with herbicide, 7-10 days before ploughing and in an interval of a least 14 days allowed before sowing.

For drilled crops, an aphicide approved for BYDV vector control can be applied at the 2-3 leaf growth stage. If low seed rates are used or the crop is early sown, an additional aphicide application may be required.

Growers should check the product label for full details.